Ref #	Hits	Search Query	DBs	Default Operator	Plurals	Time Stamp
L1	530868	poly(Gln)	US-PGPUB; USPAT; EPO; JPO; DERWENT	OR	ON	2005/04/26 09:57
L2	4475	I1 same filament	US-PGPUB; USPAT; EPO; JPO; DERWENT	OR	ON	2005/04/26 09:55
L3	13	I2 same aggregat?	US-PGPUB; USPAT; EPO; JPO; DERWENT	OR	ON	2005/04/26 09:56
L4	3	(CAG adj1 repeat) same filament	US-PGPUB; USPAT; EPO; JPO; DERWENT	OR	ON	2005/04/26 10:08
L5	12	polyglutamine same filament	US-PGPUB; USPAT; EPO; JPO; DERWENT	OR	ON	2005/04/26 10:12
L6	49	polyglutamine same aggregat?	US-PGPUB; USPAT; EPO; JPO; DERWENT	OR	ON	2005/04/26 10:12
L7	2	polyglutamine same aggregat? same diameter	US-PGPUB; USPAT; EPO; JPO; DERWENT	OR	ON	2005/04/26 10:13
L8	41831	(aggregat? or filament) same diameter	US-PGPUB; USPAT; EPO; JPO; DERWENT	OR	ON	2005/04/26 10:13
وا	4	16 and 18	US-PGPUB; USPAT; EPO; JPO; DERWENT	OR	ON	2005/04/26 10:13

FILE 'MEDICONF' ENTERED AT 10:43:14 ON 26 APR 2005 COPYRIGHT (c) 2005 FAIRBASE Datenbank GmbH, Hannover, Germany

FILE 'PASCAL' ENTERED AT 10:43:14 ON 26 APR 2005
Any reproduction or dissemination in part or in full,
by means of any process and on any support whatsoever
is prohibited without the prior written agreement of INIST-CNRS.
COPYRIGHT (C) 2005 INIST-CNRS. All rights reserved.

```
=> hayashi y/au
L37
           236 FILE AGRICOLA
L38
           734 FILE BIOTECHNO
           179 FILE CONFSCI
L39
           11 FILE HEALSAFE
L40
'AU' IS NOT A VALID FIELD CODE
L41
             0 FILE IMSDRUGCONF
           649 FILE LIFESCI
L42
'AU' IS NOT A VALID FIELD CODE
             0 FILE MEDICONF
L43
L44
          2394 FILE PASCAL
```

TOTAL FOR ALL FILES

L45 4203 HAYASHI Y/AU

=> 145 and filamentous L46 0 FILE AGRICOLA L47 1 FILE BIOTECHNO L48 0 FILE CONFSCI L49 0 FILE HEALSAFE L50 0 FILE IMSDRUGCONF 2 FILE LIFESCI L51 L52 0 FILE MEDICONF L53 2 FILE PASCAL

TOTAL FOR ALL FILES

L54 5 L45 AND FILAMENTOUS

=> dup rem
ENTER L# LIST OR (END):154

DUPLICATE IS NOT AVAILABLE IN 'IMSDRUGCONF, MEDICONF'.
ANSWERS FROM THESE FILES WILL BE CONSIDERED UNIQUE
PROCESSING COMPLETED FOR L54
L55 4 DUP REM L54 (1 DUPLICATE REMOVED)

=> d 155 ibib abs total

L55 ANSWER 1 OF 4 PASCAL COPYRIGHT 2005 INIST-CNRS. ALL RIGHTS RESERVED. on

STN

ACCESSION NUMBER: 1999-0008186 PASCAL

COPYRIGHT NOTICE: Copyright .COPYRGT. 1999 INIST-CNRS. All rights

reserved.

TITLE (IN ENGLISH): Hereditary dentatorubral-pallidoluysian atrophy:

detection of widespread ubiquitinated neuronal and

glial intranuclear inclusions in the brain

AUTHOR: HAYASHI Y.; KAKITA A.; YAMADA M.; KOIDE R.;

IGARASHI S.; TAKANO H.; IKEUCHI T.; WAKABAYASHI K.;

EGAWA S.; TSUJI S.; TAKAHASHI H.

CORPORATE SOURCE: Department of Pathology, Brain Research Institute,

Niigata University, 1-757 Asahimachi, Niigata 951-8585, Japan; Department of Neurology, Brain Research Institute, Niigata University, Niigata, Japan; Brain Disease Research Center, Brain Research

Institute, Niigata University, Niigata, Japan SOURCE:

Acta neuropathologica, (1998), 96(6), 547-552, 31

refs.

ISSN: 0001-6322 CODEN: ANPTAL

DOCUMENT TYPE: Journal BIBLIOGRAPHIC LEVEL: Analytic

COUNTRY: Germany, Federal Republic of

LANGUAGE: English

AVAILABILITY: INIST-9757, 354000071701460010

ΔN 1999-0008186 PASCAL

CP Copyright .COPYRGT. 1999 INIST-CNRS. All rights reserved. AB We examined the brains and spinal cords of seven patients with

clinicopathologically and genetically confirmed hereditary dentatorubral-pallidoluysian atrophy (DRPLA) using an antibody against ubiquitin, and found small, round immunoreactive intranuclear inclusions in both neurons and glial cells in various brain regions. Ubiquitinated neuronal intranuclear inclusions (uNIIs) were consistently found in the striatum, the pontine nuclei, the inferior olivary complex, the cerebellar cortex and the dentate nucleus. Ubiquitinated glial intranuclear inclusions (uGIIs) were found less frequently than uNIIs. Most of the inclusion-bearing nuclei were of an astrocytic nature. Immunostaining with an antibody against DRPLA protein revealed similar immunoreactive neuronal and glial intranuclear inclusions, but in much smaller in numbers compared with uNIIs and uGIIs. Electron microscopy

showed that such inclusions were composed of granular and filamentous structures. These findings strongly suggest that, in DRPLA, the occurrence of uNIIs and uGIIs is directly related to the causative gene abnormality (an expanded CAG repeat encoding polyglutamine), that neurons are affected much more widely than previously recognized and that glial cells are also involved in the

disease process.

ANSWER 2 OF 4 PASCAL COPYRIGHT 2005 INIST-CNRS. ALL RIGHTS RESERVED. on L55

STN

ACCESSION NUMBER: 1998-0237652 PASCAL

COPYRIGHT NOTICE: Copyright . COPYRGT. 1998 INIST-CNRS. All rights

reserved.

TITLE (IN ENGLISH): Hereditary dentatorubral-pallidoluysian atrophy :

ubiquitinated filamentous inclusions in the

cerebellar dentate nucleus neurons

AUTHOR: HAYASHI Y.; KAKITA A.; YAMADA M.; EGAWA S.;

OYANAGI S.; NAITO H.; TSUJI S.; TAKAHASHI H.

CORPORATE SOURCE: Department of Pathology, Brain Research Institute,

Niigata University, 1-757 Asahimachi, Niigata 951-8585, Japan; Department of Neurology, Brain Research Institute, Niigata University, 1-757

Asahimachi, Niigata 951-8585, Japan; Nagaoka Ryoikuen,

Fukazawa-cho, Nagaoka, Japan; Matsuhama Hospital,

Matsuhama-cho, Niigata, Japan

SOURCE: Acta neuropathologica, (1998), 95(5), 479-482, 25

refs.

ISSN: 0001-6322 CODEN: ANPTAL

DOCUMENT TYPE: Journal

BIBLIOGRAPHIC LEVEL: Analytic

COUNTRY: Germany, Federal Republic of

LANGUAGE: English

AVAILABILITY: INIST-9757, 354000075470040060

PASCAL ΑN 1998-0237652

Copyright .COPYRGT. 1998 INIST-CNRS. All rights reserved. CP

AB We examined the cerebellar dentate nucleus (CDN) in 16 patients with hereditary dentatorubral-pallidoluysian atrophy (DRPLA), one of the neurodegenerative diseases caused by expansion of a CAG repeat encoding a polyglutamine tract in the disease protein. In all patients, some CDN

neurons were found to contain ubiquitinated filamentous inclusions in their cytoplasm. On hematoxylin and eosin preparations, these filamentous inclusions were eosinophilic, basophilic or amphophilic, and were often found in areas of pale cytoplasm. Electron microscopy revealed that they consisted of bundles of filaments that were somewhat thicker than neurofilaments. These features of the present inclusions were indistinguishable from those of skein-like inclusions (SLI) previously described in the lower motor neurons in sporadic amyotrophic lateral sclerosis. We conclude that SLI can also occur in the CDN in DRPLA and believe that they reflect a characteristic pathological process in this disease.

ANSWER 3 OF 4 BIOTECHNO COPYRIGHT 2005 Elsevier Science B.V. on STN L55

DUPLICATE

ACCESSION NUMBER: 1998:28082456 BIOTECHNO

TITLE: Suppression of aggregate formation and apoptosis by

transglutaminase inhibitors in cells expressing

truncated DRPLA protein with an expanded polyglutamine

stretch

AUTHOR: Igarashi S.; Koide R.; Shimohata T.; Yamada M.; Hayashi Y.; Takano H.; Date H.; Oyake M.; Sato

T.; Sato A.; Egawa S.; Ikeuchi T.; Tanaka H.; Nakano

R.; Tanaka K.; Hozumi I.; Inuzuka T.; Takahashi H.;

Tsuji S.

CORPORATE SOURCE: S. Tsuji, Department of Neurology, Niigata University,

1-757 Asahimachi, Niigata 951, Japan.

E-mail: tsuji@cc.niigata-u.ac.jp

SOURCE: Nature Genetics, (1998), 18/2 (111-117), 38

reference(s)

CODEN: NGENEC ISSN: 1061-4036

DOCUMENT TYPE: Journal; Article

COUNTRY: United States

LANGUAGE: English SUMMARY LANGUAGE: English

1998:28082456 BIOTECHNO AB To elucidate the molecular mechanisms whereby expanded polyglutamine stretches elicit a gain of toxic function, we expressed full-length and truncated DRPLA (dentatorubral-pallidoluysian atrophy) cDNAs with or without expanded CAG repeats in COS-7 cells. We found that truncated DRPLA proteins containing an expanded polyglutamine stretch form

filamentous peri- and intranuclear aggregates and undergo apoptosis. The apoptotic cell death was partially suppressed by the transglutaminase inhibitors cystamine and monodansyl cadaverine (but not putrescine), suggesting involvement of a transglutaminase reaction and providing a potential basis for the development of therapeutic measures for CAG-repeat expansion diseases.

L55 ANSWER 4 OF 4 LIFESCI COPYRIGHT 2005 CSA on STN

ACCESSION NUMBER: 87:74453 LIFESCI

TITLE: Ultrastructure of cementum formation on partially formed

teeth in dogs.

Hayashi, Y.

CORPORATE SOURCE: Dep. Conserv. Dent., Fac. Dent., Kyushu Univ. 61, Maidashi

3-1-1, Higashi-ku, Fukuoka 812, Japan

SOURCE: ACTA ANAT., (1987) vol. 129, no. 4, pp. 279-288.

DOCUMENT TYPE: Journal

FILE SEGMENT:

LANGUAGE: English SUMMARY LANGUAGE: English

Cementum crystals and matrix vesicles on the root surface of partially

formed teeth in dogs were examined with a transmission electron

microscope. Fine filamentous crystals were observed in the

cementum calcifying fronts. The running pattern was mainly parallel to the

root surface in the apical region and perpendicular to the root surface in lateral and coronal regions. Matrix vesicles were observed at the apical half of the periodontium, but not observed at the coronal region. These findings suggest that the parallel-arranged cementum would become the light-microscopic lamellar type and the perpendicular one the light-microscopic dense-line structure when fully developed. Moreover, cementum formation occurs due to two kinds of mechanisms: participation of matrix vesicles and secondary calcification (= additional cementogenesis).